

SINGULAR COMPONENTS OF SPECTRAL MEASURES FOR ERGODIC JACOBI MATRICES

C. A. MARX

ABSTRACT. For ergodic 1d Jacobi operators we prove that the random singular components of any spectral measure are a.s. mutually disjoint as long as one restricts to the set of positive Lyapunov exponent. In the context of extended Harper's equation this yields the first rigorous proof of the Thouless' formula for the Lyapunov exponent in the dual regions.

1. INTRODUCTION

Given two sequences

$$(1.1) \quad (a_n) \in l^\infty(\mathbb{Z}; \mathbb{C}) , |a_n| > 0 \ \forall n \in \mathbb{Z} ,$$

$$(1.2) \quad (b_n) \in l^\infty(\mathbb{Z}; \mathbb{R}) ,$$

we define

$$(1.3) \quad (H\psi)_n := b_n\psi_n + a_n\psi_{n+1} + \overline{a_{n-1}}\psi_{n-1} ,$$

a bounded self adjoint operator on $\mathcal{H} := l^2(\mathbb{Z}; \mathbb{C})$.

Operators of the form (1.3) are known as 1 d - Jacobi operators (or Jacobi matrices), the most prominent and well studied case being Schrödinger operators where $(a_n) = (1)$.

We note that since (1.3) is unitarily equivalent to a Jacobi operator with $a_n > 0$, $\forall n$, one could assume all $a_n > 0$ right from the start. In some situations it may however be more convenient to retain the original complex form, which is why we do not apply this unitary here.

In this article, we shall be interested in an ergodic family of Jacobi operators, $(H_\omega)_{\omega \in \Omega}$, induced by random variables a, b on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$,

$$(1.4) \quad a : \Omega \rightarrow \mathbb{C}, \ b : \Omega \rightarrow \mathbb{R} ,$$

$$(1.5) \quad |\mathbb{E}_\omega \log|a|| < \infty ,$$

and an ergodic invertible map T on Ω generating the random sequences $(a_n(\omega))$ and $(b_n(\omega))$ according to

$$(1.6) \quad a_n(\omega) := a(T^n\omega) , \ b_n(\omega) = b(T^n\omega) .$$

Remark 1.1. In particular, (1.6) together with (1.5) imply that $a_n(\omega) \neq 0$, $\forall n \in \mathbb{Z}$, $\mathbb{P}$ -a.s.

The work was supported by NSF Grant DMS - 0601081 and BSF, grant 2006483.

We emphasize that in (1.3) we do not require a *positive* lower bound for the sequence (a_n) . From a dynamical point of view, fixing $z \in \mathbb{C}$, this manifests itself in almost singularity of the 1-step transfer matrices

$$(1.7) \quad B_n = \frac{1}{a_n} \begin{pmatrix} b_n - z & -\overline{a_{n-1}} \\ a_n & 0 \end{pmatrix}.$$

Here, the phrase “almost singular” refers to the matrix $a_n B_n$ whose determinant may come arbitrarily close to zero. For the ergodic counterpart in (1.5) we even allow for the matrix $a_n B_n(\omega)$ to be singular, however since $a \in L^1$ this only happens on a set of zero probability. A more detailed discussion of singular Jacobi cocycles may be found in [11].

As the main result of this article, we prove the following characterization of the singular(pp+sc)- spectrum for ergodic Jacobi matrices:

Theorem 1.2. *Let H_ω be an ergodic family of Jacobi operators satisfying (1.5) and (1.6), and let $d\mu_{\omega,s}$ be the singular component of some spectral measure for H_ω . Then, for $\mathbb{P}$ -a.e. ω and ω' , $d\mu_{\omega,s}$ and $d\mu_{\omega',s}$ are mutually singular on the set of positive Lyapunov exponent, $\{E : L(E) > 0\}$.*

Theorem 1.2 generalizes a result in [6] stated for the Schrödinger case. We note that this theorem will be proven using Kotani theory [1, 4], which usually gives results relating to the absolutely continuous component of the spectrum. It is also noteworthy that mutual singularity of the singular component of spectral measures is reminiscent of an analogous property exhibited by rank 1 perturbations of bounded s.a. operators (see e.g. [16]).

We organize the paper as follows. Sec. 2 is devoted to the proof of Theorem 1.2. As a crucial first step, we prove that energies with positive Lyapunov exponent (LE) allow for two solutions u_+, u_- of $H_\omega \psi = E\psi$ which are $\mathbb{P}$ -a.s. non-zero and l^2 (and hence necessarily exponentially decaying) at respectively $+\infty$ and $-\infty$ (Theorem 2.4).

Consequently we obtain that given an energy E with positive LE, the probability for E to be in “the natural” supporting set (defined in (2.19)) of the singular component of *any* spectral measure is zero (Theorem 2.6). This result is known for Schrödinger operators [4].

While following the basic strategy laid out in [4], our proof even when applied to the Schrödinger case differs in a number of technical details. In particular, Theorem 2.4 strengthens the respective statement for Schrödinger operators given in Theorem 6.5 of [4], by removing a set of zero probability, thus simplifying subsequent arguments (for details see remark 2.5).

Finally, Theorem 1.2 follows as a corollary of Theorem 2.6.

Sec. 3 is devoted to an application of Theorem 1.2 which really served as motivation for the results of Sec. 2. Ergodic Jacobi operators with singular transfer matrices arise naturally in a tight binding description of solids subject to an external magnetic field.

A prominent example is extended Harper's model [12, 9, 7, 18], associated with the following almost-periodic operator $H_{\bar{\lambda}, \theta; \alpha}$ defined on $l^2(\mathbb{Z})$:

$$(1.8) \quad (H_{\bar{\lambda}, \theta; \alpha} \psi)_k := v(\theta + \alpha k) \psi_k + c(\theta + \alpha k) \psi_{k+1} + \bar{c}(\theta + \alpha(k-1)) \psi_{k-1} ,$$

where $\bar{\lambda} = (\lambda_1, \lambda_2, \lambda_3)$ and

$$(1.9) \quad c_{\bar{\lambda}}(x) := \lambda_3 e^{-2\pi i(x + \frac{\alpha}{2})} + \lambda_2 + \lambda_1 e^{2\pi i(x + \frac{\alpha}{2})} , \quad v(x) := 2 \cos(2\pi x) ,$$

are analytic functions on $\mathbb{R}/\mathbb{Z}$.

The operator in (1.8) describes the influence of a transversal magnetic field to the motion of an electron in a 2-dimensional crystal layer. Here, α is a fixed irrational number associated with the magnetic flux through the unit cell and $\theta \in [0, 1)$ is the (random) quasi-momentum usually referred to as “random phase”.

The parameter triple $(\lambda_1, \lambda_2, \lambda_3)$ (*coupling*) models the lattice geometry as well as interactions between the lattice sites allowing for both nearest (λ_2) and next-nearest neighbor (λ_1 and λ_3) interactions. Without loss of generality one may assume $0 \leq \lambda_2$, $0 \leq \lambda_1 + \lambda_3$ and at least one of $\lambda_1, \lambda_2, \lambda_3$ to be positive [9]. We note that the quasi-periodic operator in (1.8) specializes to the famous almost Mathieu operator (in physics literature also known as *Harper's model*) upon setting $\lambda_1 = \lambda_3 = 0$.

Even though some attempts, both rigorous [12, 9] and heuristic [7, 18], have been made, so far little is known about extended Harper's model. As an important ingredient for the spectral analysis it is desirable to obtain the Lyapunov exponent (on the spectrum) as a function of the coupling constants.

Theorem 1.2 enables us to prove a formula of the Lyapunov exponent in the so called “dual regions” of parameter space (Theorem 3.1). A precise meaning of the word “duality” is given at the beginning of Sec. 3. We mention that Theorem 3.1 provides a rigorous proof of results obtained in physics literature by Thouless [7].

Finally, we mention that the missing link to a complete understanding of the Lyapunov exponent, the so called “self dual” region of coupling constants, a priori cannot be approached via duality. Computing the Lyapunov exponent in the self dual region therefore constitutes an important open problem which so far has escaped treatment even on a heuristic level in physics literature [7]. The development of these a priori non-duality based methods will be the subject of future work.

Acknowledgement. I would like to thank Svetlana Jitomirskaya for valuable discussions during the preparation of this manuscript. I would also like to thank the anonymous referee for insightful remarks leading to improvement of our presentation.

2. PROOF OF THE MAIN THEOREM

We start with the following basic result valid for *any* ergodic family of bounded self adjoint operators:

Theorem 2.1. *For fixed $d \in \mathbb{N}$, let H_ω be any ergodic family of bounded self adjoint operators on $l^2(\mathbb{Z}^d)$. Assume that for $\mathbb{P}$ -a.e. ω , the eigenvalues of H_ω have finite multiplicity, whenever H_ω has non-empty point spectrum. Then, fixing $E \in \mathbb{R}$, $\mathbb{P}\{\omega : E \text{ is eigenvalue for } H_\omega\} = 0$.*

Proof. Let $B \in \mathbb{R}$ be any fixed Borel set and denote by $P_B(H_\omega)$ the spectral projection onto B . The claim follows since by ergodicity $\dim \text{Ran} P_B(H_\omega)$ is constant $\mathbb{P}$ -a.s with value either ∞ or 0 [14, 3]. $\square$

In order to relate the spectral properties of the family H_ω to its dynamics given in terms of the Lyapunov exponent, we shall consider solutions $u(z)$ in $\mathbb{C}^\mathbb{Z}$ of the second order difference equation $Hu = zu$, where $z \in \mathbb{C}$ is fixed. For two such solutions u and v (same z), define the Wronskian

$$(2.1) \quad W_n[u, v] := u_n v_{n+1} - u_{n+1} v_n, n \in \mathbb{Z}.$$

It is easy to see that the Wronskian of two solutions satisfies the following conservation law

$$(2.2) \quad \overline{a_{n-1}} W_{n-1}[u, v] = a_n W_n[u, v], n \in \mathbb{Z}.$$

As an immediate consequence of (2.2) we obtain

Proposition 2.1. *Any eigenvalue of a Jacobi operator has multiplicity one.*

We note that Proposition 2.1 uses $a_n \neq 0, \forall n \in \mathbb{Z}$. In particular, in the ergodic setup, remark 1.1 implies that Proposition 2.1 holds $\mathbb{P}$ -a.s. Simplicity of the point spectrum of Jacobi operators shall be crucial for Sec. 3.

Sometimes it is useful to relate the solutions of the Schrödinger equation for the double-sided and the respective one-sided operator $H_+ := P_+ H P_+$, where P_+ is a projection onto the positive half line.

For $z \in \mathbb{C}$ fixed, a solution ψ to $H\psi = z\psi$ is a solution of the respective (positive) half-line equation if and only if it satisfies the boundary condition $\psi_0 = 0$. In particular this implies that Proposition 2.1 also applies for eigenvalues of H^+ .

More generally, for any $n \in \mathbb{Z}$ one can consider the positive n -one-sided operator $H_+^n := P^{+n} H P^{+n}$ with P^{+n} the projection onto $\{m > n\}$. For convenience, we allow $n = -\infty$ in which case we identify H_+^n with the double-sided operator H . Similarly, one defines the negative n -one-sided operator, $H_-^n := P^{-n} H P^{-n}$, $n \in \mathbb{Z}$, where P^{-n} is the projection onto $\{m < n\}$. Again, we adopt the convention $H = H_-^n$ for $n = +\infty$. To simplify notation we write $H_\pm := H_\pm^0$.

For the ergodic setup, Proposition 2.1 and Theorem 2.1 imply the following statement which will be key for the proof of Theorem 1.2:

Proposition 2.2. *Let H_ω be a family of ergodic Jacobi operators, and let $E \in \mathbb{R}$ be fixed. Then the probability for E to be an eigenvalue of some (positive or negative) n -one-sided operator (including $n = \pm\infty$) is zero.*

For $z \in \mathbb{H}^+$, let $\psi_+(z)$ be a solution to the equation $H\psi = z\psi$ which is l^2 at $+\infty$. Self adjointness of the half-line operator implies existence of $\psi_+(z)$ as well as $\psi_{+,n}(z) \neq 0$, $\forall n$. Using (2.2), we conclude uniqueness of $\psi_+(z)$ up to a multiplicative constant. Similarly, we can find a solution $\psi_-(z)$ which is l^2 at $-\infty$.

It is well known that the Green's function of H can be expressed in terms of $\psi_{\pm}(z)$ by

$$(2.3) \quad G(n, m; z) := \langle \delta_n, G(z) \delta_m \rangle = \frac{\psi_{-, \min\{n, m\}}(z) \psi_{+, \max\{n, m\}}(z)}{a_m W_m[\psi_-(z), \psi_+(z)]},$$

for $z \in \mathbb{C} \setminus \mathbb{R}$. Here, $\{\delta_n, n \in \mathbb{Z}\}$ denotes the standard basis in $l^2(\mathbb{Z})$.

Finally, for $n \in \mathbb{Z}$ we define the m -functions,

$$(2.4) \quad m_{+,n}(z) := -\frac{\psi_{+,n+1}(z)}{a_n \psi_{+,n}(z)}, \quad m_{-,n}(z) := -\frac{\psi_{-,n-1}(z)}{a_{n-1} \psi_{-,n}(z)},$$

where we set $m_{\pm}(z) := m_{\pm,0}(z)$. It is clear that in the ergodic case we have $m_{\pm,n}(\omega) = m_{\pm}(T^n \omega)$, $n \in \mathbb{Z}$. For simplicity of notation, we will suppress the z -dependence of the m -functions whenever the context permits.

Since $\psi_{\pm}$ are solutions to the Schrödinger equation, the m -functions satisfy the following Riccati-type equations

$$(2.5) \quad \begin{aligned} -\frac{1}{m_{+,n-1}} + (b_n - z) - |a_n|^2 m_{+,n} &= 0 \\ -|a_{n-1}|^2 m_{-,n} + (b_n - z) - \frac{1}{m_{-,n+1}} &= 0. \end{aligned}$$

We mention that using a relation analogous to (2.3) (see e.g. [5, 17]), the functions $m_{\pm}(z)$ relate to Borel transforms of the spectral measures $d\mu_{\pm}$ associated respectively with $H_{\pm}$ and $\delta_{\pm 1}$, i.e.

$$(2.6) \quad m_{\pm}(z) = \int \frac{d\mu_{\pm}(x)}{x - z}.$$

This in particular implies that $m_{\pm}$ are Herglotz functions, and we may use the theory of Borel transforms of measures to consider limits as $\text{Im}(z) \rightarrow 0^+$.

From here on we specifically consider the ergodic set-up. The following statement relates the Lyapunov exponent and the m -functions. It extends a result known for the Schrödinger case [15].

Proposition 2.3. *For $\text{Im } z > 0$ we have*

$$(2.7) \quad 2L(z) = \mathbb{E}_{\omega} \log \left\{ 1 + \frac{\text{Im } z}{|a_0(\omega)|^2 \text{Im } m_+(\omega, z)} \right\}.$$

Remark 2.2. We mention that a formula similar to (2.12) is given in [17] (see Eq. (5.73) therein), however with $|a_0(\omega)|^2$ in (2.3) being replaced by $a_0(\omega)$. Since a proof is omitted in [17], we correct the typo in [17] providing the details below.

Proof. Taking imaginary parts, the first equation in (2.5) implies

$$\log \left\{ |a_n(\omega)|^2 + \frac{\operatorname{Im} z}{\operatorname{Im} m_+(T^n \omega)} \right\} = \log \operatorname{Im} m_+(T^{n-1} \omega) - \log \operatorname{Im} m_+(T^n \omega) - 2 \log |m_+(T^{n-1} \omega)| .$$

Since T is measure preserving, taking expectations on both sides yields

$$(2.8) \quad \mathbb{E}_\omega \log \left\{ 1 + \frac{\operatorname{Im} z}{|a_0(\omega)|^2 \operatorname{Im} m_+(\omega)} \right\} = -2 \mathbb{E}_\omega \log \left| \frac{\psi_{+,1}(\omega)}{\psi_{+,0}(\omega)} \right| .$$

First, notice that by Birkhoff's ergodic theorem,

$$(2.9) \quad \frac{1}{n} \log \left| \frac{\psi_{+,n}(\omega)}{\psi_{+,0}(\omega)} \right| \rightarrow \mathbb{E}_\omega \log \left| \frac{\psi_{+,1}(\omega)}{\psi_{+,0}(\omega)} \right| , \quad \mathbb{P}\text{-a.e.} ,$$

as $n \rightarrow \infty$.

On the other hand $\operatorname{Im} z > 0$ implies $L(z) > 0$, hence using Oseledets' theorem (in its original $\operatorname{GL}(2, \mathbb{C})$ formulation [13]) we obtain

$$(2.10) \quad \frac{1}{n} \log \left| \frac{\psi_{+,n}(\omega)}{\psi_{+,0}(\omega)} \right| \rightarrow -L , \quad \mathbb{P}\text{-a.e.} ,$$

as $n \rightarrow \infty$. Thus, equations (2.8), (2.9), and (2.10) finally yield the claim. $\square$

Using (2.6), the monotone convergence theorem implies the existence of

$$(2.11) \quad S_\pm(\omega, E) := \lim_{\epsilon \rightarrow 0^+} \frac{\operatorname{Im} m_+(\omega, E + i\epsilon)}{\epsilon} ,$$

for any fixed $E \in \mathbb{R}$.

Thus using dominated convergence, the formula for the Lyapunov exponent given in Proposition 2.3 persists in the limit, i.e.

$$(2.12) \quad 2L(E) = \mathbb{E}_\omega \log \left\{ 1 + \frac{1}{|a_0(\omega)|^2} S_+(\omega, E)^{-1} \right\} ,$$

for any fixed $E \in \mathbb{R}$. Here, we also used that for any $E \in \mathbb{R}$, $L(E + i\epsilon)$ is continuous w.r.t. ϵ which e.g. can be obtained from the Thouless formula.

Since (2.12) contains the limit of the quantity $\frac{\operatorname{Im} m_+(\omega, z)}{\operatorname{Im} z}$, the following Lemma will prove to be of use:

Lemma 2.3. *For $\operatorname{Im} z > 0$,*

$$(2.13) \quad \frac{\operatorname{Im} m_+(\omega, z)}{\operatorname{Im} z} = \frac{1}{|a_0(\omega)|^2 |\psi_{+,0}(\omega, z)|^2} \|P_+ \psi_+(\omega, z)\|^2 .$$

Proof. Using functional calculus, Eq. (2.6) implies

$$(2.14) \quad \frac{\operatorname{Im} m_+(\omega, z)}{\operatorname{Im} z} = \sum_{n=1}^{\infty} |\langle \delta_1, (P_+ H_\omega P_+ - z)^{-1} \delta_n \rangle|^2$$

Finally, making use of expressions for the Green's function of the half line operator (see e.g. [15]), we obtain

$$(2.15) \quad |\langle \delta_1, (P_+ H_\omega P_+ - z)^{-1} \delta_n \rangle| = \left| \frac{\psi_{+,n}(\omega, z)}{a_0(\omega) \psi_{+,0}(\omega, z)} \right| ,$$

for any $n \geq 1$. $\square$

We are now in the position to characterize the set of energies with positive Lyapunov exponent in terms of the m -functions.

Theorem 2.4. *Let $E \in \mathbb{R}$ s.t. $L(E) > 0$. Then for $\mathbb{P}$ -a.e. ω , there exists a solution $u_+(\omega, E)$ with $u_{+,n}(\omega, E) \neq 0$ for all $n \in \mathbb{Z}$, which is l^2 at $+\infty$. In particular, $m_+(\omega, E + i0)$ exists, is finite and non-zero.*

Remark 2.5. In [4], the authors prove a similar statement for Schrödinger operators, which however does not rule out the case where the limiting solution $u_+(\omega, E)$ has zeros (Theorem 6.5 (b) [4]). We mention that even though the authors of [4] suspected this to only happen with zero probability, no proof was provided then.

Proof. Using (2.12), $L(E) > 0$ implies that $S_+(\omega, E) < \infty$ on a set of positive measure $\mathbb{P}$; in particular, letting $\Omega_0 := \{\omega : S_+(T^n \omega, E) < \infty \text{ for some } n \in \mathbb{Z}\}$ we obtain $\mathbb{P}(\Omega_0) > 0$. In fact, since $T^{-1}\Omega_0 = \Omega_0$, we have $\mathbb{P}(\Omega_0) = 1$ using ergodicity of T .

Let $\omega \in \Omega_0$ and $n \in \mathbb{Z}$ such that $S_+(T^n \omega, E) < \infty$. We mention that for $n = 0$, we already obtain the statement using a standard argument from the theory of Borel transforms.

Using Lemma 2.3 for $z = E + i\epsilon$ and $\epsilon > 0$,

$$(2.16) \quad \frac{\operatorname{Im} m_+(T^n \omega, z)}{\operatorname{Im} z} = \frac{1}{|a_0(T^n \omega)|^2} \|P_+ \psi_+(T^n \omega, z)\|^2 ,$$

where we define ψ_+ to satisfy the condition $\psi_{+,0}(T^n \omega, z) := 1$.

Then, by Lemma 2.3 and Banach Alaoglu ($S_+(T^n \omega, E) < \infty$), we obtain a limiting solution $u_+(\omega, E)$ of the Schrödinger equation,

$$(2.17) \quad w - \lim_{m \rightarrow \infty} \psi_+(T^m \omega, E + i\epsilon_m) =: u_+(\omega, E) ,$$

for some sequence (ϵ_m) , $\epsilon_m \searrow 0$. By construction $u_+(\omega, E)$ is l^2 at $+\infty$. Making use of (2.2), any solution of the Schrödinger equation which is l^2 at $+\infty$ is uniquely determined up to a multiplicative constant. Since by construction $u_+(n) = 1$, this limiting solution is in fact independent of the sequence (ϵ_m) , whence

$$(2.18) \quad u_+(\omega, E) = w - \lim_{\epsilon \rightarrow 0^+} \psi_+(T^n \omega, E + i\epsilon) .$$

If $u_{+,m}(\omega, E) = 0$ for some $m \in \mathbb{Z}$, $u_+(\omega, E)$ is an eigenfunction of $H_{+;\omega}^m$. By Proposition 2.2 this however happens with zero probability.

Thus, $\mathbb{P}$ -a.s. we have $u_{+,m}(\omega, E) \neq 0$, $\forall m \in \mathbb{Z}$. In particular, taking the limit in the definition of m_+ , we obtain the claim. $\square$

Following Deift and Simon [4], for $\omega \in \Omega$ we consider the following set
(2.19)

$$\mathfrak{S}_\omega := \left\{ E \in \mathbb{R} : \limsup_{\epsilon \rightarrow 0^+} \{ \operatorname{Im} G_\omega(0, 0; E + i\epsilon) + \operatorname{Im} G_\omega(1, 1; E + i\epsilon) \} = \infty \right\} .$$

It is well known that the singular component of any spectral measure for H_ω is supported on $\mathfrak{S}_\omega$ (Theorem of de la Vallée-Poussin; see also [17]).

$G(0, 0)$ and $G(1, 1)$ relate to the m-functions by

$$(2.20) \quad \begin{aligned} -G(0, 0; z)^{-1} &= |a_0|^2 m_+(z) + |a_{-1}|^2 m_-(z) + z - b_0 \\ G(1, 1; z)^{-1} &= m_+(z)^{-1} + |a_0|^2 \{ |a_{-1}|^2 m_-(z) + z - b_0 \}^{-1} . \end{aligned}$$

These relations are obtained directly from the expression for the Green's function given in (2.3).

The following statement extends the statement of Proposition 2.2 to $\mathfrak{S}_\omega$, the natural supporting set of the singular component of any spectral measure for H_ω . The respective result was first proven for Schrödinger operators by Deift and Simon [4].

Theorem 2.6. *Let $E \in \mathbb{R}$ such that $L(E) > 0$. Then, $\mathbb{P}\{\omega : E \in \mathfrak{S}_\omega\} = 0$.*

Proof. By Theorem 2.4, we know that for $\mathbb{P}$ -a.e. ω , the limits $m_\pm(\omega, E + i0)$ both exist, are finite and non-zero. Using the relations (2.20) for such ω , $E \in \mathfrak{S}_\omega$ if and only if at least one of the following happens

$$\begin{aligned} |a_0(\omega)|^2 m_+(\omega, E + i0) + |a_{-1}(\omega)|^2 m_-(\omega, E + i0) + E - b_0(\omega) &= 0 , \\ m_+(\omega, E + i0)^{-1} + |a_0(\omega)|^2 \{ |a_{-1}(\omega)|^2 m_-(\omega, E + i0) + E - b_0(\omega) \}^{-1} &= 0 . \end{aligned}$$

Using the definition of $m_\pm$, it can easily be verified that both these situations imply that the respective limiting solutions $u_\pm(\omega, E)$ constructed in the proof of Theorem 2.4 are in fact constant multiples of one another. Thus we obtain a solution which is l^2 at both $\pm\infty$, i.e. an eigenvector of H_ω . Employing Proposition 2.2, the set of such ω has zero probability, which yields the claim. $\square$

Finally we obtain Theorem 1.2, which concludes this section.

Proof of Theorem 1.2. Using the definition of the Lyapunov exponent, the set $\mathfrak{L} := \{E \in \mathbb{R} : 0 < L(E) < \infty\}$ is measurable. Consider, $\mathfrak{N} := \{(\omega, E) \in \Omega \times \mathfrak{L} : E \in \mathfrak{S}_\omega\}$. Letting $f(\omega, z) := \operatorname{Im} G_\omega(0, 0; z) + \operatorname{Im} G_\omega(1, 1; z)$, where $(\omega, z) \in \Omega \times (\mathfrak{L} + i\mathbb{R}^+)$, we note that $\mathfrak{N}$ can be written as $\mathfrak{N} = \{(\omega, E) \in \Omega \times \mathfrak{L} : \limsup_{\epsilon \rightarrow 0^+} f(\omega, E + i\epsilon) = \infty\}$. Since f is measurable, we conclude measurability of $\mathfrak{N}$.

The rest now follows using Tonelli. By Theorem 2.6, $\forall \omega' \in \Omega :$

$$\begin{aligned} 0 &= \int_{\mathfrak{L}} \mathbb{P}\{\omega : E \in \mathfrak{S}_\omega\} d\mu_{\omega'} \\ &= \int \mu_{\omega'}(\mathfrak{L} \cap \mathfrak{S}_\omega) d\mathbb{P}(\omega) , \end{aligned}$$

which implies the claim. $\square$

3. AN APPLICATION: DUALITY AND EXTENDED HARPER'S MODEL

In this section we illustrate the use of Theorem 1.2 to obtain an expression for the Lyapunov exponent in the dual regime of extended Harper's model. The model as well as its relevance has been discussed in Sec. 1.

Similar to the almost Mathieu operator, the Hamiltonian in (1.8) exhibits an intrinsic symmetry in the coupling constants under Fourier transform, known as *duality*. Heuristically speaking, duality relates certain regions in parameter space, mapping localized states to Bloch waves.

To give a precise meaning of duality, on the Hilbert space $\mathcal{H}' := L^2([0, 1) \times \mathbb{Z})$, consider the unitary operator U defined by

$$(3.1) \quad (U\phi)(\eta, m) := \sum_{n \in \mathbb{Z}} \int d\theta e^{i2\pi m\theta} e^{i2\pi n(m\alpha + \eta)} \phi(\theta, n) .$$

We note that such operator has originally been introduced by Chulaevsky-Delyon [2] and later employed in [6], both in context of the almost Mathieu operator.

For $\lambda_2 \neq 0$, let σ denote the map, $\sigma(\bar{\lambda}) := \frac{1}{\lambda_2}(\lambda_3, 1, \lambda_1)$, where as earlier $\bar{\lambda} = (\lambda_1, \lambda_2, \lambda_3)$ represents the coupling in extended Harper's model (see (1.9)).

We also define the following regions in parameter space:

- region I:** $0 \leq \lambda_1 + \lambda_3 \leq 1, 0 \leq \lambda_2 \leq 1$,
- region II:** $0 \leq \lambda_1 + \lambda_3 \leq \lambda_2, 1 \leq \lambda_2$,
- region III:** $\max\{1, \lambda_2\} \leq \lambda_1 + \lambda_3$.

This partitioning is induced by the map σ since

$$(3.2) \quad \sigma(I^\circ) = II^\circ, \quad \sigma(III) = III .$$

In this sense, the interiors of region I and II are dual, whereas region III is self-dual. For a more general discussion of duality we refer to [12].

In [9] (see Theorem 1 therein) it was proven that for Diophantine α and $\bar{\lambda} \in I^\circ$, the spectrum of $H_{\bar{\lambda}, \theta; \alpha}$ is only pure point (i.e. for a certain set $\mathbb{T}_0$ of phases θ with $|\mathbb{T}_0| = 1$). Recall, $\alpha \in \mathbb{R}$ is called Diophantine if there exists $0 < b(\alpha)$ and $1 < r(\alpha) < +\infty$ s.t. for all $j \in \mathbb{Z} \setminus \{0\}$

$$(3.3) \quad |\sin(2\pi j\alpha)| > \frac{b(\alpha)}{|j|^{r(\alpha)}} .$$

It is known that $DC(r) := \{\alpha : (3.3) \text{ holds for } r \text{ and some } b\}$ is a set of full Lebesgue measure for every $r \geq 1$.

As we will argue, Theorem 1.2 will allow to relate the regions I and II establishing the following:

Theorem 3.1. *For every Diophantine α , the LE in region II is zero on the spectrum. In region I, the Lyapunov exponent on the spectrum is positive and given*

by the formula,

$$(3.4) \quad L(\alpha; \bar{\lambda}, E) = \begin{cases} \log \left(\frac{1 + \sqrt{1 - 4\lambda_1\lambda_3}}{2\lambda_1} \right) & , \text{ if } \lambda_1 \geq \lambda_3, \lambda_2 \leq \lambda_3 + \lambda_1 , \\ \log \left(\frac{1 + \sqrt{1 - 4\lambda_1\lambda_3}}{2\lambda_3} \right) & , \text{ if } \lambda_3 \geq \lambda_1, \lambda_2 \leq \lambda_3 + \lambda_1 , \\ \log \left(\frac{1 + \sqrt{1 - 4\lambda_1\lambda_3}}{\lambda_2 + \sqrt{\lambda_2^2 - 4\lambda_1\lambda_3}} \right) & , \text{ if } \lambda_2 \geq \lambda_3 + \lambda_1 . \end{cases}$$

H in (1.8) naturally induces a bounded self adjoint operator on $\mathcal{H}'$, which shall be denoted by $H'_{\bar{\lambda};\alpha}$,

$$(3.5) \quad H'_{\bar{\lambda};\alpha} \phi := H_{\bar{\lambda};\theta,\alpha} \phi(\theta, \cdot) , \quad \forall \theta \in [0, 1) .$$

Using the definition of the unitary U , direct computation establishes the following relation between dual regions:

$$(3.6) \quad U H'_{\bar{\lambda};\alpha} U^{-1} = \lambda_2 H'_{\sigma(\bar{\lambda});\alpha} .$$

To avoid confusion, we emphasize that (3.6) shows that $H'_{\bar{\lambda};\alpha}$ and $H'_{\sigma(\bar{\lambda});\alpha}$ are unitarily equivalent and not the respective operators on the smaller space $\mathcal{H} = l^2(\mathbb{Z})$.

An important property of duality is that preserves the density of states [12]. To see this in the present formulation notice that

$$(3.7) \quad U \delta_0 = \delta_0 ,$$

where δ_0 is viewed as element of $\mathcal{H}'$, i.e. employing the embedding $\mathcal{H} \hookrightarrow \mathcal{H}'$

$$(3.8) \quad (u_n) \mapsto (\phi(\theta, n)) , \text{ where } \phi(\theta, n) = u_n \quad \forall n \in \mathbb{Z} , \theta \in [0, 1) .$$

Denote by $dn(\{H_{\bar{\lambda};\theta,\alpha}\}; E)$ the density of states of the ergodic family $\{H_{\bar{\lambda};\theta,\alpha}\}$ defined as usual by

$$(3.9) \quad \mathbb{E}_\theta \langle \delta_0, f(H_{\bar{\lambda};\theta,\alpha}) \delta_0 \rangle =: \int f(E) dn(\{H_{\bar{\lambda};\theta,\alpha}\}; E) ,$$

for f continuous and compactly supported on $\mathbb{R}$. Then, combining (3.6) and (3.7), we obtain the desired invariance of the density of states,

$$(3.10) \quad dn(\{H_{\bar{\lambda};\theta,\alpha}\}; E) = dn(\{\lambda_2 H_{\sigma(\bar{\lambda});\theta,\alpha}\}; E) = dn(\{H_{\sigma(\bar{\lambda});\theta,\alpha}\}; \lambda_2^{-1} E) .$$

We mention that the last equality in (3.10) follows from the spectral mapping theorem.

In [6] it is shown that fixing $\bar{\lambda}$ and α , it is possible to find a labeling of the eigenfunctions of $H_{\bar{\lambda};\theta,\alpha}$ that produces measurable functions w.r.t. the random phase θ .

The proof in [6] (Theorem 2.2) is given for an arbitrary bounded, ergodic Schrödinger operator on $l^2(\mathbb{Z})$, the crucial property needed being however only simplicity of the point spectrum. As shown in Proposition 2.1, the same is true

for Jacobi operators. Therefore, Theorem 2.2. in [6] extends to an ergodic family H_ω of Jacobi operators, as defined in Sec. 2.

A measurable labeling of eigenfunctions is obtained as follows: For $\omega \in \Omega$, let u be a normalized eigenvector of H_ω . Pick the unique $j \in \mathbb{Z}$ such that $|u(j)| \geq |u(k)|$, $\forall k$ and $|u(j)| > |u(k)|$, $\forall k < j$ (“left-most maximum”). We say u is attached to j . Let $N_j(\omega)$ be the number of eigenfunctions attached to j , and for $j, k \in \mathbb{Z}$ set $\Omega_{j,k} := \{\omega : N_j(\omega) \geq k\}$. On $\Omega_{j,1}$, let $(u_1(n; \omega, j))$ be the eigenfunction attached to j with maximal j th component (in magnitude) and $e_1(\omega, j)$ its associated eigenvalue. If multiple eigenfunctions attached to j have the same magnitude of their j th entry, discriminate them by choosing the one with largest eigenvalue. Note that since the point spectrum is simple, there can only be finitely many different eigenfunctions attached to j whose j th components coincide in magnitude. For further details we refer to [6].

For this labeling of eigenfunction we have (see Sec. 2 in [6]),

Theorem 3.2. *Let H_ω be an ergodic family of Jacobi operators defined as in Sec. 2, and $N_j(\omega)$, $e_l(\omega, j)$, and $(u_l(n; \omega, j))_{n \in \mathbb{Z}}$ defined as above. Then, fixing l, j and n , $N_j(\omega)$, $e_l(\omega, j)$, and $u_l(n; \omega, j)$ are measurable functions on Ω .*

The next statement shows that applying the unitary U defined in (3.1), we obtain spectral measures that are constant for a.e. phase. The proof carries over from [6] (see Theorem 3.4 therein) without any modification.

Theorem 3.3. *Fix $\bar{\lambda}$ and α Diophantine and let $(u_l(n; \theta, j))$ be the measurable labeling of eigenfunctions for $H_{\sigma(\bar{\lambda}), \theta; \alpha}$ described in Theorem 3.2. For $f(\theta) \in L^2([0, 1], d\theta)$ arbitrary and some fixed $l, j \in \mathbb{Z}$, set $\phi(\theta, n) := f(\theta)u_l(n; \theta, j)$. Denote by $d\mu_\eta$ the spectral measure for $H_{\bar{\lambda}, \theta; \alpha}$ and $\psi_\eta(n) := (U\phi)(\eta, n)$. Then, $d\mu_\eta$ is η independent for a.e. η .*

Equipped with Theorem 3.3, we are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. For a fixed Diophantine frequency α , let $\mathfrak{L}_\lambda^\pm := \{E \in \text{spec} : L(\alpha; \bar{\lambda}, E) > 0\}$. We claim that $\mathfrak{L}_\lambda^\pm = \emptyset$ for any $\bar{\lambda}$ in region II.

First consider $\bar{\lambda} \in II^\circ$ and let $d\mu_\eta$ be defined according to Theorem 3.3. Then, since $d\mu_\eta$ is constant a.e. η , Theorem 1.2 implies that in fact $d\mu_\eta$ is purely a.c. on $\mathfrak{L}_\lambda^\pm$ for a.e. η . Hence, using a well known argument by Pastur-Ishii ([8, 14]; see also the proof of Theorem 9.13 in [3]) we obtain $P_\eta^{\text{ac}}(\mathfrak{L}_\lambda^\pm) = P_\eta(\mathfrak{L}_\lambda^\pm) = 0$; here $P_\eta(B)$ and $P_\eta^{\text{ac}}(B)$ denotes the spectral projection associated with a Borel set B and its restriction to the ac subspace of H_η , respectively. In particular, in terms of the density of states this implies

$$\text{spec} = \text{supp} \{dn(\{H_{\bar{\lambda}, \theta; \alpha}\})\} \subseteq \overline{\mathbb{R} \setminus \mathfrak{L}_\lambda^\pm},$$

which by continuity of the Lyapunov exponent w.r.t. to E [11, 10] yields the claim for $\bar{\lambda} \in II^\circ$. To obtain the claim for all off region II, i.e. including the boundary, we employ the following continuity argument:

Let $\bar{\lambda}$ be an arbitrary point in ∂II and E a fixed energy in the spectrum of $H_{\bar{\lambda}}$. Using continuity of the spectrum in the Hausdorff metric ¹, there exists a sequence $(\bar{\lambda}_n)$, approximating $\bar{\lambda}$ from within II° , and a corresponding sequence of energies (E_n) , with E_n in the spectrum of $H_{\bar{\lambda}_n}$, such that $E_n \rightarrow E$. Since the Lyapunov exponent is jointly continuous at $(E, \bar{\lambda})$ for a fixed Diophantine α [11], we obtain $L(\alpha; \bar{\lambda}, E) = 0$, as claimed.

Making use of Thouless' formula and (3.10), zero Lyapunov exponent in region II immediately implies an expression for the dual region since

$$\begin{aligned}
 L(\alpha; \bar{\lambda}, E) &= - \int \log |c_{\bar{\lambda}}(x)| dx + \int \log |E - E'| dn(\{H_{\bar{\lambda}, \theta; \alpha}\}; E') \\
 (3.11) \qquad &= \int \log \left| \frac{\lambda_2 c_{\sigma(\bar{\lambda})}(x)}{c_{\bar{\lambda}}(x)} \right| dx + L(\alpha; \sigma(\bar{\lambda}), \lambda_2^{-1} E) .
 \end{aligned}$$

Here, $c_{\bar{\lambda}}$ is defined as in (1.9).

Using Jensen's formula, the integral in (3.11) can be computed explicitly [9] (see Sec. 2, p. 107),

$$(3.12) \qquad \int \log |c_{\bar{\lambda}}(x)| dx = \begin{cases} \log \lambda_3 & \text{if } \lambda_3 \geq \lambda_1 \geq 0 \\ & \text{and } \lambda_1 + \lambda_3 \geq \lambda_2 \geq 0 , \\ \log \lambda_1 & \text{if } \lambda_1 \geq \lambda_3 \geq 0 \\ & \text{and } \lambda_1 + \lambda_3 \geq \lambda_2 \geq 0 , \\ \log \left| \frac{2\lambda_1 \lambda_3}{-\lambda_2 + \sqrt{\lambda_2^2 - 4\lambda_1 \lambda_3}} \right| & \text{if } \lambda_1 + \lambda_3 \leq \lambda_2 \text{ and } \lambda_1, \lambda_3 \neq 0 , \\ \log \lambda_2 & \text{if } \lambda_1 + \lambda_3 \leq \lambda_2 , \lambda_1 \text{ or } \lambda_3 = 0 , \end{cases}$$

giving rise to the expression for the Lyapunov exponent in region I given in (3.4). $\square$

REFERENCES

1. R. Carmona, S. Kotani, *Inverse Spectral theory for random Jacobi matrices*, Journal of Stat. Phys. 46 (1987), 1091 - 1114.
2. V. Chulaevsky and F. Delyon, *Purely absolutely continuous spectrum for almost Mathieu operators*, Journal of Stat. Phys. 55 (1989), 1279 - 1284.
3. H.L. Cycon, R.G. Froese, W. Kirsch, B. Simon, *Schrödinger Operators with Application to Quantum Mechanics and Global Geometry*, Springer (1987), Heidelberg.
4. P. Deift and B. Simon, *Almost periodic Schrödinger operators, III. The absolutely continuous spectrum in one dimension*, Commun. Math. Phys. 90 (1983), 389-411.
5. F. Gesztesy and B. Simon, *m-functions and inverse spectral analysis for finite and semi-infinite Jacobi matrices*, J. d'Analyse Math. 73 (1997), 267-297.

¹It is a well known fact that given a sequence (A_n) of bounded self adjoint operators approximating a bounded operator A in *norm topology*, the spectra satisfy $\text{spec}(A_n) \rightarrow \text{spec}(A)$ in the Hausdorff metric.

6. A.Y. Gordon, S. Jitomirskaya, Y. Last and B. Simon, *Duality and singular continuous spectrum in the almost Mathieu equation*, Acta Mathematica 178 (1997), 169 - 183.
7. J. H. Han and D. J. Thouless, H. Hiramoto, M. Kohmoto, *Critical and bicritical properties of Harper's equation with next-nearest neighbor coupling*, Phys. Rev. B 50 (1994), 11365 - 11380.
8. K. Ishii, *Localization of eigenstates and transport phenomena in one-dimensional disordered systems*, Suppl. Prog. Theor. Phys. 53 (1973), 77.
9. S. Jitomirskaya, D.A. Koslover and M.S. Schulteis, *Localization for a Family of One-dimensional Quasi-periodic Operators of Magnetic Origin*, Ann. Henri Poincaré 6 (2005), 103 - 124.
10. S. Jitomirskaya, D.A. Koslover, M.S. Schulteis, *Continuity of the Lyapunov Exponent for analytic quasi-periodic cocycles*, Ergod. Th. & Dynam. Sys. 29 (2009), 1881 - 1905.
11. S. Jitomirskaya, C.A. Marx, *Continuity of the Lyapunov Exponent for analytic quasi-periodic cocycles with singularities*, Journal of Fixed Point Theory and Applications, Festschrift on occasion of the 80th birthday of Dick Palais, to appear.
12. V.A. Mandelshtam, S. Ya. Zhitomirskaya, *1D-Quasiperiodic Operators. Latent Symmetries*, Commun. Math. Phys. 139 (1991), 589 - 604.
13. V. Oseledets, *Oseledets theorem*, Scholarpedia 3(1):1846 (2008).
14. L. Pastur, *Spectral properties of disordered systems in one-body approximation*, Commun. Math. Phys. 75 (1980), 179.
15. B. Simon, *Kotani theory for one dimensional stochastic Jacobi matrices*, Commun. Math. Phys. 89 (1983), 227-234.
16. B. Simon, *Trace ideals and their applications*, 2nd edition, Amer. Math. Soc., Providence, RI, 2005.
17. G. Teschl, *Jacobi Operators and Completely Integrable Nonlinear Lattices*, Mathematical Surveys and Monographs, 72, Amer. Math. Soc., Providence 2000.
18. D. J. Thouless, *Bandwidths for a quasiperiodic tight-binding model*, Phys. Rev. B 28 (1983), 4272 - 4276.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE CA, 92717